

NASA Contractor Report 182053

ICASE Report No. 90-42

AD-A225 291

ICASE

THE CFL CONDITION FOR SPECTRAL APPROXIMATIONS
TO HYPERBOLIC INITIAL-BOUNDARY VALUE PROBLEMS

David Gottlieb
Eitan Tadmor

Contract No. NAS1-18605
June 1990

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, Virginia 23665-5225

Operated by the Universities Space Research Association

NASA

National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23665-5225

DTIC
ELECTE
JUL 18 1990
S E D

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

90 01 11 103

THE CFL CONDITION FOR SPECTRAL APPROXIMATIONS TO HYPERBOLIC INITIAL-BOUNDARY VALUE PROBLEMS¹

David Gottlieb²

Division of Applied Mathematics
Brown University
Providence, RI 02192

and

Eitan Tadmor³

School of Mathematical Sciences
Tel-Aviv University
Tel-Aviv 69978, ISRAEL

c. 100

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

ABSTRACT

We study the stability of spectral approximations to scalar hyperbolic initial-boundary value problems with variable coefficients. Time is discretized by explicit multi-level or Runge-Kutta methods of order ≤ 3 (forward Euler time differencing is included), and we study spatial discretizations by spectral and pseudospectral approximations associated with the general family of Jacobi polynomials. We prove that these fully explicit spectral approximations are stable provided their time-step, Δt , is restricted by the CFL-like condition, $\Delta t < \text{Const. } N^{-2}$, where N equals the spatial number of degrees of freedom. We give two independent proofs of this result, depending on two different choices of appropriate L^2 -weighted norms. In both approaches, the proofs hinge on a certain inverse inequality interesting for its own sake. Our result confirms the commonly held belief that the above CFL stability restriction, which is extensively used in practical implementations, guarantees the stability (and hence the convergence) of fully-explicit spectral approximations in the non-periodic case.

¹Research was supported in part by NASA Contract No. NAS1-18605 while both authors were in residence at the Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, VA 23665. The first author was also supported by the Air Force office of Scientific Research grant no. AFOSR-90-0093, by DARPA-URI contract N00014-86-k0754, and by NSF grant DMS-88-10150.

²School of Mathematical Sciences, Tel-Aviv University, Tel-Aviv 69978, ISRAEL.

³Department of Mathematics, University of Michigan, Ann Arbor, MI 48109.

1. INTRODUCTION.

We are concerned here with fully-discrete spectral and pseudospectral approximations to scalar hyperbolic equations. In this context, the spectral (and respectively, the pseudospectral) approximations consist of truncation (and respectively, collocation) of N -term spatial expansions, which are expressed in terms of general Jacobi polynomials; Chebyshev and Legendre expansions are the ones most frequently found in practice. In this paper we prove that such N -terms approximations are stable, provided their time-step, Δt , fulfills the CFL-like condition, $\Delta t \leq \text{Const.} N^{-2}$. To clarify the origin of such CFL-like conditions in our case, we recall that the Jacobi polynomials are in fact the eigenfunctions of second-order singular Sturm-Liouville problems. Our arguments show that the main reason for the above CFL limitation is the $O(N^2)$ growth of the corresponding N -eigenvalues associated with these Sturm-Liouville problems.

The paper is organized as follows. Section 2 includes a brief summary on the properties of Jacobi polynomials (and their quadrature rules) which are used throughout the paper. In Section 3 we state our main stability theorems for Forward Euler time-differencing and (pseudo-)spectral spatial differencing, for constant coefficients equations with homogeneous boundary conditions. Section 4 extends our stability results to the inhomogeneous case. In Section 5 we discuss multi-level and Runge-Kutta time differencing. Finally, in Section 6 we show how to extend our results in the presence of (positive) variable coefficients.

2. VERY SHORT GUIDE TO JACOBI POLYNOMIALS.

Jacobi polynomials, $P_k^{\alpha, \beta}$, are the eigenfunctions of the singular Sturm-Liouville problem

$$(2.1a) \quad \left((1-x^2)w(x)P_k^{(\alpha, \beta)'}(x) \right)' + \lambda_k w(x)P_k^{(\alpha, \beta)}(x) = 0, \quad -1 \leq x \leq 1,$$

with corresponding eigenvalues λ_k ,

$$(2.1b) \quad \lambda_k \equiv \lambda_k(\alpha, \beta) = k(k + \alpha + \beta + 1).$$

Different families of Jacobi polynomials are associated with different weight functions $w(x)$,

$$(2.1c) \quad w(x) \equiv w(x; \alpha, \beta) = (1-x)^\alpha (1+x)^\beta, \quad \alpha, \beta > -1.$$

In the sequel we shall frequently use several properties of the Jacobi polynomials. A brief summary of these properties is given below (consult e.g. [13]). We start with the well-known

PROPERTY 1 (Orthogonality). *We have*

$$(2.2) \quad (P_j^{(\alpha, \beta)}, P_k^{(\alpha, \beta)})_{w(x)} = 0, \quad j \neq k.$$

The derivatives of Jacobi polynomials are also Jacobi polynomials. This is evident from the following property which shows that $\{P_k^{(\alpha, \beta)'}\}_{k \geq 0}$ are orthogonal with respect to the weight $(1 - x^2)w(x) \equiv w(x; \alpha + 1, \beta + 1)$ and hence

$$(2.3) \quad P_{k+1}^{(\alpha, \beta)'} = \text{Const}_{k, \alpha, \beta} P_k^{(\alpha+1, \beta+1)}, \quad \text{Const}_{k, \alpha, \beta} = \frac{1}{2}(k + \alpha + \beta + 2).$$

PROPERTY 2 (Orthogonality of derivatives). *We have*

$$(2.4) \quad (P_j^{(\alpha, \beta)'}, P_k^{(\alpha, \beta)'})_{(1-x^2)w(x)} = 0, \quad j \neq k,$$

$$(2.5) \quad \|P_k^{(\alpha, \beta)'}\|_{(1-x^2)w(x)}^2 = \lambda_k \|P_k^{(\alpha, \beta)}\|_{w(x)}^2.$$

Indeed, (2.4) and (2.5) follow from integration by parts of (2.1) against $P_j^{(\alpha, \beta)}(x)$.

Let π_N denote the space of algebraic polynomials with degree $\leq N$. A useful consequence of the last two properties is provided by

LEMMA 2.1 (Inverse inequality). *For all $p \in \pi_N$ we have*

$$(2.6) \quad \|p'\|_{(1-x^2)w(x)} \leq \sqrt{\lambda_N} \|p\|_{w(x)}, \quad p \in \pi_N.$$

Here $w(x)$ stands for an arbitrary $w(x; \alpha, \beta)$ weight, and $\lambda_N \equiv \lambda_N(\alpha, \beta)$ is the corresponding N -th eigenvalue.

Remarks.

1. The inequality (2.6) can be viewed as the algebraic analogue of the trigonometric inverse inequality,

$$(2.7a) \quad \|p'\|_{L^2[-\pi, \pi]} \leq N \|p\|_{L^2[-\pi, \pi]}, \quad p = \text{any } N - \text{trigonometric polynomial}.$$

This should be contrasted with a similar L^2 -inverse inequality for algebraic polynomials where there is a loss of N^2 -factor for each derivative [3],

$$(2.7b) \quad \|p'\|_{L^2[-1, 1]} \leq \text{Const. } N^2 \|p\|_{L^2[-1, 1]}, \quad p = \text{any } N - \text{algebraic polynomial},$$

and this estimate, (2.7b), is sharp in view of, e.g., $p_N(x) = \sum_{k=0}^N P_k^{(-\frac{1}{2}, -\frac{1}{2})}(x)$. Thus, the use of the different weighted L^2 -norms in the algebraic case, (2.6), is essential in order to retain a loss of only $\sqrt{\lambda_N} \sim N$ -factor for each derivative.

2. The inverse inequality (2.6) can be viewed as an L^2 -weighted version of Bernstein's inequality

$$\|(1-x^2)^{\frac{1}{2}}p'(x)\|_{L^\infty[-1,1]} \leq N\|p(x)\|_{L^\infty[-1,1]}, \quad p \in \pi_N.$$

Standard interpolation arguments between this L^∞ -type estimate and the L^2 -type estimate (2.6) yield for $q \geq 2$

$$(2.7c) \quad \|(1-x^2)^{\frac{1}{2}}p'(x)\|_{L^q_{w(\cdot;\alpha,\beta)}[-1,1]} \leq \sqrt{\lambda_N(\alpha,\beta)} \cdot \|p(x)\|_{L^q_{w(\cdot;\alpha,\beta)}[-1,1]}, \quad p \in \pi_N.$$

Similar weighted L^q -type estimates apply to higher derivatives.

PROOF. Given $p(x)$ in π_N , we will use its Jacobi expansion, $p(x) = \sum_{k=0}^N a_k P_k^{(\alpha,\beta)}(x)$ and $p'(x) = \sum_{k=0}^N a_k P_k^{(\alpha,\beta)'}(x)$. Starting with the left-hand side of (2.6) and using (2.4), (2.5) and (2.2) in this order, we obtain

$$(LHS)^2 = \sum_{k=0}^N a_k^2 \|P_k^{(\alpha,\beta)'}\|_{(1-x^2)w(x)}^2 = \sum_{k=0}^N \lambda_k a_k^2 \|P_k^{(\alpha,\beta)}\|_{w(x)}^2 \leq \lambda_N (RHS)^2. \quad \square$$

We note in passing that Lemma 2.1 can be generalized to higher derivatives: successive application of (2.6) with $\omega(x) = w(x; \alpha, \beta)$ yields

$$(2.8) \quad \|p^{(k)}(x)\|_{(1-x^2)^k w(x)} \leq \prod_{j=0}^{k-1} \lambda_N(\alpha+j, \beta+j) \cdot \|p(x)\|_{w(x)}, \quad p \in \pi_N.$$

This leads us to a 'natural' definition of non-periodic Sobolev spaces equipped with finite $H_{w(x)}^s$ -norm, where,

$$(2.9) \quad \|p\|_{H_{w(x)}^s}^2 = \sum_{k=0}^s \|p^{(k)}\|_{(1-x^2)^k w(x)}^2.$$

With this in mind, we now recover a sharp inverse inequality familiar from the trigonometric setup

$$(2.10) \quad \|p\|_{H_{w(x)}^s} \leq \text{Const}_s \cdot N^s \|p\|_{w(x)}, \quad \text{Const} \sim 1 + \frac{s}{N}, \quad p \in \pi_N.$$

In the above discussion we can replace integrals by discrete summations in view of the well-known

PROPERTY 3 (Gauss quadrature rule). Let $\{q_N(x)\}_{N \geq 0}$ be a family of π_N -polynomials orthogonal w.r.t. the $\omega(x)$ -weighted L^2 inner product. Let $-1 < x_1 < x_2 \dots < x_N < 1$ be

the N zeroes of $q_N(x)$. Then there exists positive discrete weights, $\{\omega_j\}_{j=1}^N$, such that for all $p \in \pi_{2N-1}$ we have

$$(2.11) \quad \int_{-1}^1 \omega(x) p(x) dx = \sum_{j=1}^N \omega_j p(x_j), \quad p \in \pi_{2N-1}.$$

Remark. To compute the Gauss weights we set $p(x) = \frac{q_N(x)}{x - x_k}$ in (2.11). Since $p(x_j) = 0$, $j \neq k$, (2.11) yields

$$(2.12) \quad \omega_k = \frac{1}{q'_N(x_k)} \int_{-1}^1 \omega(x) \frac{q_N(x)}{x - x_k} dx, \quad 1 \leq k \leq N.$$

PROOF. We have $p(x) = t(x)q_N(x) + r(x)$ for some $t(x)$ and $r(x)$ in π_{N-1} . The choice of weights in (2.12) guarantees that (2.11) is valid for π_{N-1} -polynomials, for

$$\text{span} \left\{ \frac{q_N(x)}{x - x_k} \right\}_{1 \leq k \leq N} = \pi_{N-1}.$$

This together with our assumption that $q_N(x)$ is $L^2_{\omega(x)}$ -orthogonal to π_{N-1} imply

$$\int_{-1}^1 \omega(x) p(x) dx = \int_{-1}^1 \omega(x) r(x) dx = \sum_{j=1}^N \omega_j r(x_j) = \sum_{j=1}^N \omega_j p(x_j). \quad \square$$

EXAMPLES.

1. Gauss-Jacobi quadrature rule. By Property 1, (2.11) applies to $\{P_N^{(\alpha, \beta)}\}_{N \geq 1}$ with $\omega(x) = w(x; \alpha, \beta)$. Hence there exist $\{w_j = w_j^G(\alpha, \beta)\}_{j=1}^N$ such that

$$(2.13) \quad \int_{-1}^1 w(x) p(x) dx = \sum_{j=1}^N w_j p(x_j), \quad \text{for all } p \in \pi_{2N-1}.$$

Remark. The Gauss-Jacobi quadrature rule (2.13) can be used as a highly accurate quadrature rule for general smooth, not necessarily polynomial functions. The error incurred in such cases is governed by [4, p. 75]

$$(2.14) \quad \int_{-1}^1 w(x) f(x) dx - \sum_{j=1}^N w_j f(x_j) = \text{Const. } f^{(2N)}(\theta), \quad \text{Const.} > 0, \quad |\theta| \leq 1.$$

2. Gauss-Lobatto-Jacobi quadrature rule. By Property 2, (2.11) applies to $\{P_{N+1}^{(\alpha, \beta)}\}_{N \geq 0}$ with $\omega(x) = (1 - x^2)w(x; \alpha, \beta)$, and therefore, there exist $\{\tilde{w}_j = w_j^G(\alpha + 1, \beta + 1)\}_{j=1}^N$ such that

$$(2.15) \quad \int_{-1}^1 (1 - x^2)w(x)r(x)dx = \sum_{j=1}^N \tilde{w}_j r(x_j), \text{ for all } r \in \pi_{2N-1}.$$

This is in fact a special case of the Gauss-Lobatto-Jacobi quadrature rule which is exact for all $p \in \pi_{2N+1}$. Indeed, for all such p 's we have $p(x) = (1 - x^2)r(x) + \ell(x)$ with $r(x)$ in π_{2N-1} and a linear $\ell(x) = p(-1)\frac{1-x}{2} + p(1)\frac{1+x}{2}$. By (2.15)

$$\begin{aligned} \int_{-1}^1 w(x)p(x)dx &= \sum_{j=1}^N \tilde{w}_j r(x_j) + \int_{-1}^1 w(x)\ell(x)dx \\ &= \sum_{j=1}^N \frac{\tilde{w}_j}{1-x_j^2} p(x_j) + \int_{-1}^1 w(x)\ell(x)dx - \sum_{j=1}^N \frac{\tilde{w}_j}{1-x_j^2} \ell(x_j) = I + II + III. \end{aligned}$$

Thus, we have

$$I = \sum_{j=1}^N w_j^L p(x_j), \quad w_j^L \equiv \frac{\tilde{w}_j}{1-x_j^2} = \frac{1}{1-x_j^2} w_j^G(\alpha + 1, \beta + 1)$$

and the two expressions, $II + III$, amount to a linear combination of $p(-1)$ and $p(1)$

$$II + III = w_0^L p(x_0) + w_{N+1}^L p(x_{N+1}), \quad x_0 \equiv -1 < x_1 < \dots < x_N < 1 \equiv x_{N+1}.$$

Hence, there exist $\{w_j = w_j^L(\alpha, \beta)\}_{j=0}^{N+1}$ such that

$$(2.16) \quad \int_{-1}^1 w(x)p(x)dx = \sum_{j=0}^{N+1} w_j p(x_j), \quad \text{for all } p \in \pi_{2N+1}. \quad \square$$

Finally, we shall need some information on the behavior of the collocation points which appear on the right of (2.13) and (2.16). We have

PROPERTY 4 (Distribution of zeros). If $x_j = \cos \theta_j$ is the j -th zero of $P_N^{(\alpha, \beta)}(x)$, then [1, p. 287] $N\theta_j$ is a corresponding zero of Bessel's function, and hence

$$1 - x_j^2 = \sin^2 \theta_j \sim \text{Const}_{j_0} N^{-2} \quad \text{for } j \in J = \{1 \leq j \leq j_0, \quad N - j_0 \leq j \leq N\}.$$

Thus, the zeros of $P_N^{(\alpha, \beta)}(x)$ are accumulated within $O(N^{-2})$ neighborhood of $\{-1, +1\}$. More precise estimates, e.g., [12, p. 19] yield

$$(2.17) \quad \frac{1}{1 - x_j} \leq \frac{N^2}{2(1 + \alpha)}, \quad 1 \leq j \leq N.$$

3. FORWARD EULER WITH HOMOGENEOUS BOUNDARY CONDITIONS.

We start with the scalar constant coefficient hyperbolic equation,

$$(3.1) \quad u_t = au_x, \quad (x, t) \in [-1, 1] \times [0, \infty), \quad a > 0,$$

which is augmented with the homogeneous condition at the inflow boundary,

$$(3.2) \quad u(1, t) = 0, \quad t > 0.$$

To approximate (3.1), we use the forward Euler time differencing on the left, and either spectral or pseudospectral differencing on the right. Thus, we seek a temporal sequence of spatial π_N -polynomials, $v^m = v_N(x, t^m = m\Delta t)$, such that

$$(3.3a) \quad v_N(x, t^m + \Delta t) = v_N(x, t^m) + \Delta t \cdot v'_N(x, t^m) + \Delta t \cdot \tau(t^m)q_N(x).$$

Here, $q_N(x)$ is a π_N -polynomial which characterizes the specific (pseudo)spectral method we employ, and $\tau = \tau(t^m)$ is a free scalar multiplier to be determined by the boundary constraint

$$(3.3b) \quad v_N(x = 1, t^m) = 0.$$

We shall study the spectral-Jacobi tau methods, [8],[2], where

$$(3.4) \quad v_N(x, t^m + \Delta t) = v_N(x, t^m) + \Delta t \cdot av'_N(x, t^m) + \Delta t \cdot \tau(t^m)q_N(x), \quad q_N(x) = P_N^{(\alpha, \beta)}(x),$$

and the pseudospectral-Jacobi methods, [5],[2], which are collocated at the interior extrema of $P_{N+1}^{(\alpha, \beta)}$, i.e.,

$$(3.5) \quad v_N(x, t^m + \Delta t) = v_N(x, t^m) + \Delta t \cdot av'_N(x, t^m) + \Delta t \cdot \tau(t^m)q_N(x), \quad q_N(x) = P_{N+1}^{(\alpha, \beta)'}(x).$$

Remark. These two families of spectral and pseudospectral Jacobi methods are closely related since $P_{N+1}^{(\alpha, \beta)'}(x)$ is a scalar multiple of $P_N^{(\alpha+1, \beta+1)}(x)$, consult (2.3). We will not discuss here a different alternative to (3.5) where one collocates at the interior extrema of $P_N^{(\alpha, \beta)}(x)$ together with the downstream outflow boundary so that

$$(3.6) \quad q_N(x) = (1+x)P_N^{(\alpha, \beta)'}(x).$$

Let $-1 < x_1 < x_2 < \dots < x_N < 1$ be the N -different zeros of the forcing polynomial $q_N(x)$. The spectral approximation (3.3a) restricted to these points reads

$$(3.7a) \quad v_N(x_j, t^{m+1}) = v_N(x_j, t^m) + \Delta t \cdot av'_N(x_j, t^m), \quad 1 \leq j \leq N,$$

and is augmented with the homogeneous boundary condition

$$(3.7b) \quad v_N(1, t^m) = 0.$$

Equations (3.7a), (3.7b) furnish a complete equivalent formulation of the spectral approximation (3.3a), (3.3b). An essential ingredient in a stability theory of such approximations lies in the choice of appropriate L^2 -weighted norms

$$(3.8) \quad \|f(x)\|_{\omega}^2 = \langle f(x), f(x) \rangle, \quad \langle f(x), g(x) \rangle = \sum_{j=1}^N \omega_j f(x_j) g(x_j).$$

We make

DEFINITION 3.1 (Stability). The approximation (3.7a), (3.7b) is stable if there exist discrete weights, $\{\omega_j > 0\}_{j=1}^N$, and a constant η_0 independent of N , such that

$$(3.9) \quad \|v_N(\cdot, t)\|_{\omega} \leq \text{Const.} e^{\eta_0 t} \|v_N(\cdot, 0)\|_{\omega}.$$

The approximation (3.7a), (3.7b) is strongly stable if (3.9) holds with $\text{Const.} = 1$ and $\eta_0 \leq 0$, i.e., if

$$(3.10) \quad \|v_N(\cdot, t)\|_{\omega} \leq \|v_N(\cdot, 0)\|_{\omega}.$$

We recall that in the Jacobi-type spectral approximations (3.4) and (3.5), the nodes $\{x_j\}_{j=1}^N$ are the zeros of Jacobi polynomials associated with the Gauss and Gauss-Lobatto quadrature rules. We use

$$(3.11) \quad \Delta x_{\min} = \min(1 + x_1, 1 - x_N)$$

to measure the minimal gridsize associated with these Gauss nodes. Our choice of discrete weights $\{\omega_j\}_{j=1}^N$ for the stability of the spectral and pseudospectral-Jacobi methods (3.4), (3.5) will be specified later on; these weights are related (but not equal) to the corresponding Gauss weights $\{w_j\}_{j=1}^N$ indicated earlier.

With this in mind we have

THEOREM 3.1 (Stability of the spectral and pseudospectral Jacobi methods). *Consider the spectral approximations (3.7a), (3.7b), associated with the Jacobi-tau method (3.4) or the pseudospectral-Jacobi method (3.5). There exists a positive constant $\eta_0 \equiv \eta_0(\alpha, \beta) > 0$ independent of N , such that if the following CFL condition holds*

$$(3.12) \quad \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \leq \eta_0,$$

then the approximation (3.7a), (3.7b) is strongly stable and the following estimate is fulfilled

$$(3.13) \quad \|v_N(\cdot, t)\|_\omega \leq e^{-\eta_0 \alpha t} \|v_N(\cdot, 0)\|_\omega.$$

Notes.

1. The choice of L^2 -weighted norms. Theorem 3.1 deals with the stability of both the spectral-tau methods associated with Jacobi polynomials $P_N^{(\alpha, \beta)}(x)$, $\alpha, \beta \in (-1, 0)$, and with the closely related pseudospectral methods associated with $P_{N+1}^{(\alpha, \beta)'}(x)$, $\alpha, \beta \in (-1, 0)$. In each case we give two different stability proofs which are based on two different choices of discrete L^2 -weighted norms; these discrete weights $\{\omega_j\}_{j=1}^N$ are given by

$$(3.14a) \quad \omega_j = \frac{1+x_j}{1-x_j} w_j, \quad \text{where } \{w_j\}_{j=1}^N = \text{Gauss - Jacobi weights in (2.13),}$$

$$(3.14b) \quad \omega_j = (1+x_j)w_j, \quad \text{where } \{w_j\}_{j=1}^N = \text{Gauss - Lobatto Jacobi weights in (2.16).}$$

2. The CFL condition. The CFL condition (3.12) places an $O(N^{-2})$ stability restriction on the time-step Δt , and this stability restriction involves two factors. First, since we expand our solution in terms of the eigenfunctions of the Sturm-Liouville problem (2.1), the CFL condition involves the corresponding $(N-1)$ -th eigenvalue

$$(3.15a) \quad \lambda_{N-1} \equiv \lambda_{N-1}(\alpha, \beta) < N^2, \quad \alpha, \beta \in (-1, 1).$$

Second, the spectral Jacobi approximation (3.7a) is collocated at Gauss nodes which are accumulated within $O(N^{-2})$ neighborhoods near the boundaries, i.e., by (2.17),

$$(3.15b) \quad \frac{1}{\Delta x_{\min}} \leq \frac{N^2}{2(1+\alpha)}.$$

Thus, in view of (3.15a), (3.15b), the CFL condition (3.12) boils down to

$$(3.16) \quad \Delta t \cdot \alpha l^2 \leq \text{Const.}, \quad \text{Const.} = \eta_0 \frac{1+\alpha}{2+\alpha} > 0.$$

3. The choice of a stability norm. The stability statement asserted in Theorem 3.1 is formulated in terms of discrete semi-norms, $\|\cdot\|_\omega$, which are ω -weighted by either (3.14a) or (3.14b). We note that $\|\cdot\|_\omega$ are in fact well-defined norms on the space of π_N -polynomials

satisfying the vanishing boundary condition (3.7b), i.e., corresponding to (3.14a) or (3.14b) we have

$$(3.17a) \quad \|v_N(\cdot, t)\|_\omega = \int_{-1}^1 w(x) \frac{1+x}{1-x} v_N^2(x, t) dx, \quad v_N(1, t) = 0,$$

$$(3.17b) \quad \|v_N(\cdot, t)\|_\omega = \int_{-1}^1 w(x)(1+x) v_N^2(x, t) dx, \quad v_N(1, t) = 0.$$

Moreover, in view of (3.15b), one may convert the stability statement (3.13) into the usual L_2 -type stability estimate at the expense of possible algebraic growth which reads

$$(3.18) \quad \|v_N(\cdot, t)\|_{w(x)} \leq \frac{N^2}{1+\alpha} e^{-\eta_0 a t} \|v_N(\cdot, 0)\|_{w(x)}, \quad \|v_N(\cdot, t)\|_{w(x)}^2 = \int_{-1}^1 w(x) v_N^2(x, t) dx.$$

4. Exponential time decay. Let us integrate by parts the differential equation (3.1) against $(1+x)u$. Thanks to the homogeneous boundary condition (3.2) we find

$$(3.19) \quad \frac{d}{dt} \int_{-1}^1 (1+x) u^2(x, t) dx \leq -\frac{a}{2} \int_{-1}^1 (1+x) u^2(x, t) dx,$$

and therefore,

$$(3.20) \quad \|u(\cdot, t)\|_{1+x} \leq e^{-\frac{1}{4} a t} \|u(\cdot, 0)\|_{1+x}.$$

This estimate corresponds to the special case of the stability statement (3.13) for the spectral-Legendre tau method ($\alpha = \beta = 0$) weighted by (3.14b). The exponential time decay indicated in (3.20) and more generally in (3.13), is due to the special choice of ω -weighted stability norms. The weights $\{w_j\}_{j=1}^N$ in (3.14a), (3.14b) involve the essential factors $1+x_j$ or $\frac{1+x_j}{1-x_j}$ which amplify the inflow boundary values in comparison to the outflow ones. Since in the current homogeneous case, vanishing inflow data is propagating into the domain, this results in the exponential time decay indicated in (3.20) and likewise in the stability statement (3.13).

5. The inflow problem. A stability statement similar to Theorem 3.1 is valid in the inflow case where $a < 0$. Assume that the CFL condition (3.12) holds with $\eta_0 = \eta_0(\beta, \alpha)$, then (3.13) follows with discrete weights $\omega_j = \frac{1-x_j}{1+x_j} w_j$ or $\omega_j = (1-x_j) w_j$.

The rest of this section is devoted to the proof of Theorem 3.1 according to the various cases outlined in the four lemmata below. We start with

LEMMA 3.2 (Stability of the spectral-tau method). *Consider the spectral-Jacobi-tau method (3.4). Then Theorem 3.1 holds with*

$$(3.21a) \quad \omega_j = \frac{1+x_j}{1-x_j} w_j, \quad \{w_j = w_j^G(\alpha, \beta)\}_{j=1}^N = \text{Gauss - Jacobi weights},$$

$$(3.21b) \quad \eta_0 \equiv \eta_0(\alpha, \beta) = \begin{cases} \frac{1}{2}(1+\beta), & \alpha + \beta \leq 0, \\ \frac{1}{2}(1-\alpha), & \alpha + \beta \geq 0. \end{cases} \quad \alpha, \beta \in (-1, 1).$$

PROOF. Squaring of (3.7a) yields

$$(3.22) \quad \begin{aligned} \|v_N(\cdot, t^{m+1})\|_\omega^2 &= \|v_N(\cdot, t^m)\|_\omega^2 + \\ &+ 2\Delta t \cdot a < v_N(\cdot, t^m), v'_N(\cdot, t^m) > + (\Delta t \cdot a)^2 \|v'_N(\cdot, t^m)\|_\omega^2 = \\ &= \|v_N(\cdot, t^m)\|_\omega^2 + 2\Delta t \cdot a I + (\Delta t \cdot a)^2 II, \end{aligned}$$

and we turn to estimate the two expressions, I and II, on the right of (3.22).

First let us note that since the π_N -polynomial $v_N(x, t^m)$ vanishes at the inflow boundary, (3.3b), we have

$$(3.23) \quad v_N(x, t^m) = (1-x)p(x) \quad \text{for some } p(x) \equiv p_{N-1}(x) \in \pi_{N-1}.$$

Also, a straightforward computation shows that

$$(3.24) \quad \left(w(x) \frac{1+x}{1-x} \right)' (1-x)^2 = [(\beta - \alpha + 2) - (\beta + \alpha)x] w(x) \geq 4\eta_0 w(x), \quad |x| \leq 1,$$

where $\eta_0 = \eta_0(\alpha, \beta)$ is given in (3.21b).

Now, since $\frac{1+x}{1-x} v_N(x, t^m) v'_N(x, t^m) \in \pi_{2N-1}$, Gauss quadrature rule (2.11) implies

$$I \equiv \sum_{j=1}^N w_j \frac{1+x_j}{1-x_j} v_N(x_j, t^m) v'_N(x_j, t^m) = \int_{-1}^1 w(x) \frac{1+x}{1-x} v_N(x, t^m) v'_N(x, t^m) dx.$$

We integrate by parts the RHS of I, substitute $v_N(x, t^m) = (1-x)p(x)$ from (3.23), and in view of (3.24) we obtain

$$(3.25) \quad I = -\frac{1}{2} \int_{-1}^1 \left(w(x) \frac{1+x}{1-x} \right)' (1-x)^2 p^2(x) dx \leq -2\eta_0 \|p\|_{w(x)}^2.$$

Next, let us consider the second expression, II, on the right of (3.22). As before, we substitute $v_N(x, t^m) = (1-x)p(x)$ from (3.23) and obtain

$$II \equiv \|v'_N(\cdot, t^m)\|_\omega^2 = \sum_{j=1}^N w_j \frac{1+x_j}{1-x_j} [(1-x_j)p'(x_j) - p(x_j)]^2 \leq$$

$$\leq 2 \sum_{j=1}^N w_j (1 - x_j^2) (p'(x_j))^2 + 2 \sum_{j=1}^N w_j \frac{1 + x_j}{1 - x_j} p^2(x_j) = II_1 + II_2.$$

Since $(1 - x^2)(p'(x))^2 \in \pi_{2N-2}$, the Gauss quadrature rule (2.13) followed by the inverse inequality (2.6) implies

$$II_1 \equiv 2 \sum_{j=1}^N w_j (1 - x_j^2) (p'(x_j))^2 = 2 \|p'\|_{(1-x^2)w(x)}^2 \leq 2\lambda_{N-1} \|p\|_{w(x)}^2, \quad p \in \pi_N.$$

This together with the obvious upper-bound

$$II_2 \equiv 2 \sum_{j=1}^N w_j \frac{1 + x_j}{1 - x_j} p^2(x_j) \leq \frac{4}{\Delta x_{\min}} \|p\|_{w(x)}^2,$$

gives us

$$(3.26) \quad II \leq \left(2\lambda_{N-1} + \frac{4}{\Delta x_{\min}} \right) \|p\|_{w(x)}^2.$$

Equipped with (3.25) and (3.26) we return to (3.22) to find

$$(3.27) \quad \|v_N(\cdot, t^{m+1})\|_{\omega}^2 \leq \|v_N(\cdot, t^m)\|_{\omega}^2 - 2\Delta t \cdot a \left[2\eta_0 - \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \right] \|p\|_{w(x)}^2.$$

The CFL condition (3.21b) implies that the squared brackets on the right are nonnegative,

$$(3.28) \quad \left[2\eta_0 - \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \right] \geq \eta_0 > 0,$$

and hence strong stability.

In fact, one more application of the Gauss quadrature rule yields

$$(3.29) \quad \begin{aligned} \|p\|_{w(x)}^2 &= \sum_{j=1}^N w_j p^2(x_j) = \sum_{j=1}^N w_j \frac{v_N^2(x_j, t^m)}{(1-x_j)^2} \geq \\ &\geq \sum_{j=1}^N w_j \frac{1+x_j}{1-x_j} v_N^2(x_j, t^m) = \|v_N(\cdot, t^m)\|_{\omega}^2. \end{aligned}$$

The inequalities (3.29), (3.28) together with (3.27) imply

$$(3.30) \quad \|v_N(\cdot, t^{m+1})\|_{\omega}^2 \leq (1 - 2\eta_0 \Delta t \cdot a) \|v_N(\cdot, t^m)\|_{\omega}^2$$

and the result (3.13) follows. \square

Next, we take advantage of the rather general setup we employed in Lemma 3.2. Specifically, since $P_{N+1}^{(\alpha, \beta)'}$ is proportional to $P_N^{(\alpha+1, \beta+1)}$, consult (2.3), we may use Lemma 3.2 with $\eta_0(\alpha, \beta)$ replaced by $\eta_0(\alpha+1, \beta+1)$ to conclude:

LEMMA 3.3 (Stability of the pseudospectral method). *Consider the pseudospectral-Jacobi method (3.5). Then Theorem 3.1 holds with*

$$(3.31a) \quad \omega_j = \frac{1+x_j}{1-x_j} w_j, \quad \{w_j = w_j^G(\alpha+1, \beta+1)\}_{j=1}^N = \text{Gauss - Jacobi weights},$$

$$(3.31b) \quad \eta_0 \equiv \eta_0(\alpha, \beta) = -\frac{\alpha}{2} > 0, \quad \alpha, \beta \in (-1, 0).$$

As mentioned before, alternative proofs of Theorem 3.1 are possible. For example, following [6, Theorem 5.1] one may employ a stable norm weighted by $\omega_j = (1+x_j)w_j$ instead of the $\omega_j = \frac{1+x_j}{1-x_j}w_j$ weights used in (3.21a), (3.31a). We begin with

LEMMA 3.4 (Stability of the spectral-tau method revisited). *Consider the spectral-Jacobi tau method (3.4). Then Theorem 3.1 holds with*

$$(3.32a) \quad \omega_j = (1+x_j)w_j, \quad \{w_j = w_j^G(\alpha, \beta)\}_{j=1}^N = \text{Gauss - Jacobi weights}$$

$$(3.32b) \quad \eta_0 = \eta_0(\alpha, \beta) = \begin{cases} -\frac{\alpha}{2}, & \alpha + \beta + 1 \geq 0, \\ \frac{1}{2}(1 - \beta), & \alpha + \beta + 1 \leq 0, \end{cases} \quad \alpha, \beta \in (-1, 0).$$

PROOF. We square (3.7a) – this time using the inner product weighted by (3.32a),

$$(3.33) \quad \begin{aligned} \|v_N(\cdot, t^{m+1})\|_\omega^2 &= \|v_N(\cdot, t)\|_\omega^2 + 2\Delta t \cdot a < v_N(\cdot, t^m), v'_N(\cdot, t^n) > + \\ &+ (\Delta t \cdot a)^2 \|v'_N(\cdot, t^m)\|_\omega^2 = \\ &= \|v_N(\cdot, t^m)\|_\omega^2 + 2\Delta t \cdot a I + (\Delta t \cdot a)^2 II, \end{aligned}$$

and as before we have to estimate the two expressions on the right of (3.33).

The first expression, I, involves discrete summation of $f(x) = (1+x)v_N(x, t^m)v'_N(x, t^m)$, and since $f(x)$ is a π_{2N} -polynomial, its N -nodes Gaussian sum is not an exact integral. The essential observation here is that $f^{(2N)} \equiv \text{Const.} > 0$ in this case, and the error estimate (2.14) tells us that the Gauss quadrature rule is upper bounded by

$$(3.34) \quad I \equiv \sum_{j=1}^N w_j(1+x_j)v_N(x_j, t^m)v'_N(x_j, t^m) < \int_{-1}^1 w(x)(1+x)v_N(x, t^n)v'_N(x, t^m)dx.$$

Let us recall that the homogeneous inflow boundary condition (3.7b) implies

$$(3.35) \quad v_N(x, t^m) = (1-x)p(x), \text{ for some } p(x) \equiv p_{N-1}(x) \in \pi_{N-1}.$$

Also, a straightforward computation shows

$$(3.36) \quad (w(x)(1+x))'(1-x) = [(\beta - \alpha + 1) - (\alpha + \beta + 1)x]w(x) \geq 4\eta_0 w(x), \quad |x| \leq 1$$

where $\eta_0 = \eta_0(\alpha, \beta)$ is given in (3.32b).

We integrate by parts the RHS of I, substitute (3.35), and in view of (3.36) we obtain

$$(3.37) \quad I = -\frac{1}{2} \int_{-1}^1 (w(x)(1+x))'(1-x)^2 p^2(x) dx \leq -2\eta_0 \|p\|_{(1-x)w(x)}^2.$$

Next, let us consider the second expression, II, on the right of (3.33). As before, we substitute $v_N(x, t^m) = (1-x)p(x)$ from (3.35) and Gauss quadrature yields

$$(3.38) \quad \begin{aligned} II &= \sum_{j=1}^N w_j(1+x_j)[(1-x_j)p'(x_j) - p(x_j)]^2 = \\ &= \int_{-1}^1 w(x)(1-x^2)(1-x)(p'(x))^2 - 2 \int_{-1}^1 w(x)(1-x^2)p(x)p'(x) dx \\ &\quad + \int_{-1}^1 w(x)(1+x)p^2(x) dx = II_1 + II_2 + II_3. \end{aligned}$$

The inverse inequality (2.6) with weight $\omega(x) = (1-x)w(x)$ implies

$$II_1 = \|p'\|_{(1-x^2)(1-x)w(x)}^2 \leq \lambda_{N-1} \|p\|_{(1-x)w(x)}^2, \quad \lambda_{N-1} = \lambda_{N-1}(\alpha + 1, \beta),$$

and integration by parts of II_2 gives

$$\begin{aligned} II_2 + II_3 &= \int_{-1}^1 [(w(x)(1-x^2))' + w(x)(1+x)]p^2(x) dx \\ &\leq 2\|p^2\|_{w(x)} = 2 \sum_{j=1}^N w_j \frac{1-x_j}{1-x_j} p^2(x_j) \leq \frac{2}{\Delta x_{\min}} \|p\|_{(1-x)w(x)}^2. \end{aligned}$$

Consequently we have

$$(3.39) \quad II \leq \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \|p\|_{(1-x)w(x)}^2.$$

Equipped with (3.37) and (3.39) we return to (3.33) to find

$$(3.40) \quad \|v_N(\cdot, t^{m+1})\|_{\omega}^2 \leq \|v_N(\cdot, t^m)\|_{\omega}^2 - 2\Delta t \cdot a \left[2\eta_0 - \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \right] \|p\|_{(1-x)w(x)}^2$$

and the result follows along the lines of Lemma 3.2, consult (3.27). \square

Lemma 3.4 does not cover the pseudospectral Jacobi methods, since by (2.6) the corresponding Jacobi parameters $\alpha + 1, \beta + 1 \notin (-1, 0)$. However, the proof of Lemma 3.4 can be carried out in the pseudospectral case if we replace the Gauss quadrature rule by the Gauss-Lobatto one. We omit the almost identical details (which are outlined for the variable coefficients case in Theorem 6.2 below) and state

LEMMA 3.5 (Stability of the pseudospectral method revisited). *Consider the pseudospectral Jacobi method (3.5). Then Theorem 3.1 holds with*

$$(3.41a) \quad \omega_j = (1 + x_j)w_j, \quad \{w_j = w_j^L(\alpha, \beta)\}_{j=1}^N = \text{Gauss - Lobatto - Jacobi weights},$$

$$(3.41b) \quad \eta_0 = \eta_0(\alpha, \beta) = \begin{cases} -\frac{\alpha}{2}, & \alpha + \beta + 1 \geq 0, \\ \frac{1}{2}(1 - \beta), & \alpha + \beta + 1 \leq 0. \end{cases} \quad \alpha, \beta \in (-1, 0).$$

Remark. The stability asserted in Lemma 3.5 is stated in terms of the discrete seminorm $\|v_N(\cdot, t)\|_\omega^2 = \sum_{j=1}^N w_j(1 + x_j)v_N^2(x_j, t)$ weighted by the interior Gauss-Lobatto weights $\{w_j\}_{j=1}^N$. However, taking into account the homogeneous boundary condition (3.7b) and the exactness of Gauss-Lobatto quadrature for π_{2N+1} polynomials, this amounts to

$$\|v_N(\cdot, t)\|_\omega^2 = \sum_{j=0}^{N+1} w_j(1 + x_j)v_N^2(x_j, t) = \int_{-1}^1 w(x)(1 + x)v_N^2(x, t)dx.$$

4. FORWARD EULER WITH INHOMOGENEOUS INITIAL-BOUNDARY CONDITIONS.

We consider the inhomogeneous scalar hyperbolic equation

$$(4.1) \quad u_t = au_x + F(x, t), \quad (x, t) \in [-1, 1] \times [0, \infty), \quad a > 0,$$

which is augmented with inhomogeneous data prescribed at the inflow boundary

$$(4.2) \quad u(1, t) = g(t), \quad t > 0.$$

Using the forward Euler time differencing, the spectral approximation of (4.1) reads, at the N -interior zeros of $q_N(x)$,

$$(4.3a) \quad v_N(x_j, t^{m+1}) = v_N(x_j, t^m) + \Delta t \cdot av'_N(x_j, t^m) + \Delta t F(x_j, t^m), \quad q_N(x_j) = 0,$$

and is augmented with the boundary condition

$$(4.3b) \quad v_N(1, t^m) = g(t^m).$$

In this section, we study the stability of (4.3), (4.4) in the two cases of

$$(4.4a) \quad \text{Spectral - Jacobi tau method: } q_N(x) = P_N^{(\alpha, \beta)}(x), \quad \alpha, \beta \in (-1, 0),$$

and the closely related

$$(4.4b) \quad \text{pseudospectral - Jacobi method : } q_N(x) = P_{N+1}^{(\alpha, \beta)'}(x), \quad \alpha, \beta \in (-1, 0).$$

To deal with the inhomogeneity of the boundary condition (4.3b), we employ a device introduced in [6, Section 5]. Namely we consider the π_N -polynomial

$$(4.5) \quad V_N(x, t) = v_N(x, t) - \frac{q_N(x)}{q_N(1)} g(t).$$

If we set

$$(4.6) \quad \tilde{F}(x, t) = F(x, t) + a \frac{q'_N(x)}{q_N(1)} g(t)$$

then $V_N(x, t)$ satisfies the inhomogeneous equation

$$(4.7a) \quad V_N(x_j, t^{m+1}) = V_N(x_j, t^m) + \Delta t \cdot a V'_N(x_j, t^m) + \Delta t \tilde{F}(x_j, t^m),$$

which is now augmented by the homogeneous boundary condition

$$(4.7b) \quad V_N(1, t^m) = 0.$$

Theorem 3.1 together with Duhammel's principle provides us with a priori estimate of $\|V_N(\cdot, t)\|_\omega$ in terms of the initial and the inhomogeneous data, $\|V_N(\cdot, 0)\|_\omega$ and $\|\tilde{F}(\cdot, t)\|_\omega$. Namely, if the CFL condition (3.12) holds, then we have

$$(4.8) \quad \|V_N(\cdot, t)\|_\omega \leq e^{-\eta_0 a t} \|V_N(\cdot, 0)\|_\omega + \sum_{0 < t^m \leq t} \Delta t \cdot e^{-\eta_0 a(t-t^m)} \|\tilde{F}(\cdot, t^m)\|_\omega.$$

Since the discrete norm $\|\cdot\|_\omega$ is supported at the interior zeros of $q_N(x)$ where $V_N(x_j, t) = v_N(x_j, t)$ we conclude

THEOREM 4.1 (Stability of the spectral tau and pseudospectral Jacobi methods). *Consider the spectral approximation (3.3a), (3.3b) associated with the Jacobi-tau method (4.4a) or the pseudospectral-Jacobi method (4.4b). There exists a positive constant $\eta_0 = \eta_0(\alpha, \beta) > 0$ independent of N , such that if the following CFL condition holds (consult (3.12)),*

$$(4.9) \quad \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \leq \eta_0$$

then the approximation (3.3a), (3.3b) satisfies the stability estimate

$$(4.10) \quad \|v_N(\cdot, t)\|_\omega \leq e^{-\eta_0 a t} \|v_N(\cdot, 0)\|_\omega + \sum_{0 < t^m \leq t} \Delta t \cdot e^{-\eta_0 a(t-t^m)} \left[\|F(\cdot, t^m)\|_\omega + a \frac{\|q'_N(\cdot)\|_\omega}{|q_N(1)|} |g(t^m)| \right].$$

Theorem 4.1 provides us with a priori stability estimate in terms of the initial data, $v_N(\cdot, 0)$, the inhomogeneous data, $F(\cdot, t)$, and the boundary data $g(t)$. The dependence on the boundary data involves the factor of $\frac{\|q'_N(\cdot)\|_\omega}{|q_N(1)|}$, which grows linearly with N , as shown by

LEMMA 4.2. *Let $\{w_j\}_{j=1}^N$ be the discrete weights given by either (3.32a) or (3.41a). Then there exists a constant independent of N such that*

$$(4.11) \quad \frac{\|q'_N(\cdot)\|_\omega}{|q_N(1)|} \leq \text{Const. } N.$$

Remark. The stability estimate (4.10) together with (4.11) imply

$$(4.12) \quad \|v_N(\cdot, t)\|_\omega \leq e^{-\eta_0 a t} \|v_N(\cdot, 0)\|_\omega + \sum_{0 < t^m \leq t} \Delta t \cdot e^{-\eta_0 a(t-t^m)} [\|F(\cdot, t^m)\|_\omega + \text{Const. } N \Delta t \cdot e^{-\eta_0 a(t-t^m)} |g(t^m)|].$$

An inequality similar to (4.12) is encountered in the stability study of finite-difference approximations to mixed initial-boundary hyperbolic systems [9]. We note in passing that the stability estimate (4.12) together with the usual consistency requirement guarantee the spectrally accurate convergence of the spectral approximation; consult [7] for the semi-discrete case.

PROOF. We consider, for example, the spectral-tau method associated with $q_N(x) = P_N^{(\alpha, \beta)}(x)$ and with discrete weights $\omega_j = (1 + x_j)w_j^G(\alpha, \beta)$. Using Gauss and Gauss-Lobatto quadrature rules (2.13) and (2.16) in this order, we obtain

$$\|q'_N(\cdot)\|_\omega^2 = \int_{-1}^1 w(x)(1+x)(P_N^{(\alpha, \beta)'}(x))^2 dx = 2w_{N+1}^L(\alpha, \beta) |P_N^{(\alpha, \beta)'}(1)|^2,$$

and (4.11) follows in view of

$$2w_{N+1}^L(\alpha, \beta) \left| \frac{P_N^{(\alpha, \beta)'}(1)}{P_N^{(\alpha, \beta)}(1)} \right|^2 \leq \text{Const. } N^2.$$

Similar arguments (which are omitted) apply to the pseudospectral approximation associated with $q_N(x) = P_N^{(\alpha, \beta)'}(x)$ and with discrete weights $\omega_j = (1 + x_j)w_j^L(\alpha, \beta)$.

5. MULTI-LEVEL AND RUNGE-KUTTA TIME DIFFERENCING.

In the previous sections we proved the stability of spectral approximations which are combined with the first order accurate forward Euler time differencing. In this section we

extend our stability result for certain second and third order accurate multi-level and Runge-Kutta time differencing, which were studied in [10], [11].

To this end we view our π_N -approximate solution at time level t , $v(\cdot, t)$, as an $(N + 1)$ -dimensional column vector which is uniquely realized at the Gauss collocation nodes $(v(x_1, t), \dots, v(x_N, t), v(1, t))$.

The forward Euler time differencing with homogeneous boundary conditions (3.4), (3.5), reads

$$(5.1a) \quad v(t^m + \Delta t) = [I + \Delta t \cdot aL]v(t^m), \quad a > 0,$$

where L is an $(N + 1) \times (N + 1)$ matrix which accounts for the spatial spectral differencing together with the homogeneous boundary conditions

$$(5.1b) \quad Lv(t^m) = (v'(x_1, t^m), \dots, v'(x_N, t^m), 0).$$

Remark. The construction of a spectral differentiation matrix L can be accomplished in one of two ways. One possibility is carried out in the physical space, by exact differentiation of the unique π_N -interpolant which assumes the gridvalues $v(x_1, t), \dots, v(x_N, t), v(1, t)$ at the corresponding Gauss nodes. This leads to full $(N + 1) \times (N + 1)$ differentiation matrices L , which are recorded for example in [2]. Spectral differencing in the physical space is then carried out by a matrix-vector multiplication at the expense of $O(N^2)$ operations. Alternatively, spectral differentiation can be accomplished in the transformed space. In the particular case of Chebyshev (pseudo-)spectral method, this leads to a factorization of the corresponding differentiation matrix L , whose matrix-vector multiplication can be carried out efficiently by FFT requiring $O(N \log N)$ operations, consult [8], [2].

Theorem 3.1 tells us that if the CFL condition (3.12) holds, i.e., if

$$(5.2) \quad \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \leq \eta_0,$$

then $I + \Delta t \cdot aL$ is bounded in the ω -weighted induced operator norm,

$$(5.3) \quad \|I + \Delta t \cdot aL\|_{\omega} \leq e^{-\eta_0 a \cdot \Delta t}.$$

Let us consider an $(s + 2)$ -level time differencing method of the form

$$(5.4) \quad v(t^m + \Delta t) = \sum_{k=0}^s \theta_k [I + c_k \Delta t \cdot aL] v(t^{m-k}), \quad c_k \geq 0, \quad \theta_k \geq 0, \quad \sum_{k=0}^s \theta_k = 1.$$

In this case, $v(t^m + \Delta t)$ is given by a convex combination of stable forward Euler differencing and we conclude

COROLLARY 5.1 (Multi-level time differencing). Assume that the following CFL condition holds,

$$(5.5) \quad \Delta t \cdot a \left(\lambda_{N-1} + \frac{2}{\Delta x_{\min}} \right) \leq \frac{\eta_0(\alpha, \beta)}{c_k}, \quad c_k \geq 0, \quad k = 0, 1, \dots, s.$$

Then the spectral approximation (5.4) is strongly stable and the following estimate holds

$$(5.6) \quad \|v_N(\cdot, t)\|_{\omega} \leq e^{-\eta_* \Delta t} \|v_N(\cdot, 0)\|_{\omega}, \quad \eta_* = \min_k \frac{\eta_0}{c_k} > 0.$$

In [10], second and third order accurate multi-level time differencing methods of the positive type (5.4) were constructed. They take the particularly simple form

$$(5.7) \quad v(t^m + \Delta t) = \theta[I + c_0 \Delta t \cdot aL]v(t^m) + (1 - \theta)[I + c_s \Delta t \cdot aL]v(t^{m-s}),$$

with positive coefficients given in Table 1.

Second-order time differencing	θ	c_0	c_s
4-level method ($s = 2$)	$\frac{3}{4}$	2	0
5-level method ($s = 3$)	$\frac{8}{9}$	$\frac{3}{2}$	0
Third-order time differencing			
5-level method ($s = 3$)	$\frac{16}{27}$	3	$\frac{12}{11}$
6-level method ($s = 4$)	$\frac{25}{32}$	2	$\frac{10}{7}$
7-level method ($s = 5$)	$\frac{108}{125}$	$\frac{5}{3}$	$\frac{30}{17}$

Table 1. Multi-level methods.

Similar arguments apply for Runge-Kutta time differencing methods. In this case the resulting positive type Runge-Kutta methods take the form

$$(5.8a) \quad v^{(1)}(t^{m+1}) = [I + \Delta t \cdot aL]v(t^m),$$

$$(5.8b) \quad v^{(k)}(t^{m+1}) = \theta_k v(t^m) + (1 - \theta_k)[I + \Delta t aL]v^{(k-1)}(t^{m+1}), \quad k = 2, \dots, s,$$

$$(5.8c) \quad v(t^{m+1}) = v^{(s)}(t^{m+1}).$$

We state

COROLLARY 5.2 (Runge-Kutta time differencing). *Assume that the CFL condition (3.12) holds. Then the spectral approximation (5.8a-c) with $0 \leq \theta_k < 1$ is strongly stable and the stability estimate (3.13) holds.*

In Table 2 we quote the preferred second and third-order choices of [11].

Second order time differencing	θ_2	θ_3
Two-step modified Euler ($s = 2$)	$\frac{1}{2}$	—
Third order time differencing		
Three-step method ($s = 3$)	$\frac{3}{4}$	$\frac{1}{3}$

Table 2. Runge-Kutta methods

Remarks.

1. The above results can be extended to include nonhomogeneous data as well. We omit the details.

2. In [10], [11], higher order accurate (> 3) multi-level and Runge-Kutta time differencing schemes were constructed. In the present context (of constant coefficient spectral approximations), they amount to convex combinations of the stable forward Euler differencing $I + \Delta t \cdot aL$ and its adjoint $I - \Delta t \cdot aL$. The above argument does not cover these cases, however, since in our case the stability of $I \pm \Delta t \cdot aL$ is measured by different weighted norms.

6. SCALAR EQUATIONS WITH VARIABLE COEFFICIENTS.

We begin with

EPILOGUE. When dealing with finite-difference approximations which are locally supported, i.e., finite difference schemes whose stencil occupies a finite number of neighboring grid cells each of which is of size Δx , then one encounters the hyperbolic CFL stability restriction

$$(6.1) \quad \frac{\Delta t}{\Delta x} |a| \leq \text{Const.}$$

With this in mind, it is tempting to provide a heuristic justification for the stability of spectral methods, by arguing that a CFL stability restriction similar to (6.1) should hold. Namely, when Δx is replaced by the minimal gridsize, $\Delta x_{\min} = \min_j |x_{j+1} - x_j| = O(N^{-2})$, then (6.1) leads us to

$$(6.2) \quad \Delta t \cdot |a| N^2 \leq \text{Const.}$$

Although the final conclusion is correct (consult (3.16)), it is important to realize that this "handwaving" argument is not well-founded in the case of spectral methods. Indeed, since the spectral stencil occupies the whole interval $(-1,1)$, spectral methods do not lend themselves to the stability analysis of locally supported finite-difference approximations. Of course, by the same token this explains the existence of unconditionally stable fully implicit (and hence globally supported) finite difference approximations.

As noted earlier, our stability proof (in Theorem 3.1) shows that the CFL condition (6.2) is related to the following two points:

#1. The size of the corresponding Sturm-Liouville eigenvalues, $\lambda_{N-1} = O(N^2)$.

#2. The minimal gridsize, $\frac{1}{\Delta x_{\min}} = O(N^2)$.

The second point seems to support the fact that Δx_{\min} plays an essential role in the CFL stability restriction for the global spectral methods, as predicted by the local heuristic argument outlined above. To clarify this issue we study in this section the stability of spectral approximations to scalar hyperbolic equations with variable coefficients. The principle *raison d'être* which motivates our present study, is to show that our stability analysis in the constant coefficients case is versatile enough to deal with certain variable coefficients problems. We begin with the particular example introduced in [8]

$$(6.3) \quad u_t = -xu_x, \quad (x, t) \in [-1, 1] \times [0, \infty).$$

We shall show that the CFL stability restriction in this case is related to the $O(N^2)$ -size of the Sturm-Liouville eigenvalues (point #1 above), but otherwise it is independent of the minimal gridsize mentioned in point #2 above.

Observe that (6.3) requires no augmenting boundary conditions, since both boundaries, $x = \pm 1$, are outflow ones. Consequently, the various π_N -spectral approximations of (6.3) with forward Euler time differencing, read

$$(6.4) \quad v_N(x, t^m + \Delta t) = v_N(x, t^m) - \Delta t \cdot x v'_N(x, t^m).$$

We have

THEOREM 6.1 (Stability). *Assume that the following CFL condition holds*

$$(6.5) \quad \Delta t \cdot \lambda_N \leq 1, \quad \lambda_N = N(N+1).$$

Then the spectral approximation (6.4) is strongly stable and the following estimate is fulfilled

$$(6.6) \quad \|v_N(\cdot, t)\|_{1-x^2} \leq \|v_N(\cdot, 0)\|_{1-x^2}.$$

PROOF. Squaring (6.4) yields

$$(6.7) \quad \begin{aligned} \|v_N(\cdot, t^{m+1})\|_{1-x^2}^2 &= \|v_N(\cdot, t^m)\|_{1-x^2}^2 + \\ &-2\Delta t(v_N(\cdot, t^m), xv'_N(\cdot, t^m))_{1-x^2} + (\Delta t)^2 \|xv'_N(\cdot, t^m)\|_{1-x^2}^2 \\ &= \|v_N(\cdot, t^m)\|_{1-x^2}^2 + 2\Delta t \cdot I + (\Delta t)^2 \cdot II. \end{aligned}$$

Integration by parts shows that the first expression, I, is given by

$$(6.8) \quad I \equiv \frac{1}{2} \int_{-1}^1 (x(1-x^2))' v_N^2(x, t^m) dx = \frac{1}{2} \|v_N(\cdot, t^m)\|_{1-x^2}^2 - \int_{-1}^1 x^2 v_N^2(x, t^m) dx.$$

Next, we write $xv'_N \equiv (xv_N)' - v_N$, and by the inverse inequality (2.6) the second expression, II, does not exceed

$$(6.9) \quad II = \|(xv_N(x, t^m))'\|_{1-x^2}^2 - 2 \int_{-1}^1 x^2 v_N^2(x, t^m) dx \leq (\lambda_N - 2) \int_{-1}^1 x^2 v_N^2(x, t^m) dx.$$

Inserting (6.8) and (6.9) into (6.7) we end up with

$$(6.10) \quad \|v_N(\cdot, t^{m+1})\|_{1-x^2}^2 \leq (1 + \Delta t) \cdot \|v_N(\cdot, t^m)\|_{1-x^2}^2 + \Delta t \cdot [(\lambda_N - 2)\Delta t - 2] \cdot \int_{-1}^1 x^2 v_N^2(x, t) dx.$$

The CFL condition (6.5) tells us that the contribution of the second term is negative, for

$$\Delta t [(\lambda_N - 2)\Delta t - 2] \cdot \int_{-1}^1 x^2 v_N^2(x, t^m) dx \leq -\Delta t \|v(\cdot, t^m)\|_{1-x^2}^2$$

and the strong stability estimate (6.6) now follows. \square

We now turn to discuss scalar hyperbolic equations with positive variable coefficients

$$(6.11) \quad u_t = a(x)u_x, \quad 0 < a(x) < a_\infty, \quad (x, t) \in [-1, 1] \times [0, \infty),$$

which are augmented with homogeneous conditions at the inflow boundary

$$(6.12) \quad u(1, t) = 0.$$

We consider the pseudospectral-Jacobi method collocated at the N -interior zeros of $P_{N+1}^{(\alpha, \beta)'}(x)$. Using forward Euler time differencing, the resulting approximation reads

$$(6.13a) \quad v_N(x_j, t^{m+1}) = v_N(x_j, t^m) + \Delta t \cdot a(x_j) v_N'(x_j, t^m), \quad P_{N+1}^{(\alpha, \beta)'}(x_j) = 0,$$

together with the boundary condition

$$(6.13b) \quad v_N(1, t^m) = 0.$$

Arguing along the lines of Theorem 3.1, we have

THEOREM 6.2 (Stability of the pseudospectral Jacobi method with variable coefficients).

Consider the pseudospectral-Jacobi approximation (6.13a), (6.13b). There exists a constant $\eta_0 \equiv \eta_0(\alpha, \beta)$,

$$(6.14a) \quad \eta_0 \equiv \eta_0(\alpha, \beta) = \begin{cases} -\frac{\alpha}{2}, & \alpha + \beta + 1 \geq 0, \\ \frac{1}{2}(1 - \beta), & \alpha + \beta + 1 \leq 0, \end{cases} \quad \alpha, \beta \in (-1, 0),$$

such that if the following CFL condition holds

$$(6.14b) \quad \Delta t \left(a_\infty \lambda_{N-1} + 2 \max_{1 \leq j \leq N} \frac{a(x_j)}{1 - x_j} \right) \leq \eta_0,$$

then the approximation (6.13a), (6.13b) is strongly stable, i.e., there exist discrete weights

$$(6.15a) \quad \omega_j = (1 + x_j) \frac{w_j}{a(x_j)}, \quad \{w_j = w_j^L(\alpha, \beta)\}_{j=1}^N = \text{Gauss - Lobatto weights},$$

such that

$$(6.15b) \quad \|v_N(\cdot, t)\|_\omega \leq \|v_N(\cdot, 0)\|_\omega.$$

PROOF. We divide (6.13a) by $\sqrt{a(x_j)}$,

$$\frac{1}{\sqrt{a(x_j)}} v_N(x_j, t^{m+1}) = \frac{1}{\sqrt{a(x_j)}} v_N(x_j, t^m) + \Delta t \cdot \sqrt{a(x_j)} \cdot v_N'(x_j, t^m),$$

and, proceeding as before, we square both sides to obtain

$$(6.16) \quad \begin{aligned} \|v_N(\cdot, t^{m+1})\|_\omega^2 &= \|v_N(\cdot, t^m)\|_\omega^2 + \\ &+ 2\Delta t \langle v_N(\cdot, t^m), v_N'(\cdot, t^m) \rangle + (\Delta t)^2 \|a(\cdot) v_N'(\cdot, t^m)\|_\omega^2 \\ &= \|v_N(\cdot, t^m)\|_\omega^2 + 2\Delta t \cdot I + (\Delta t)^2 \cdot II. \end{aligned}$$

The first expression, I, involves discrete summation of the π_{2N} -polynomial $f(x) = (1 + x)v_N(x, t^m)v'_N(x, t^m)$ and since $f(\pm 1) = 0$ (in view of (6.13b)), the $(N + 1)$ -nodes Gauss-Lobatto quadrature rule yields

$$I \equiv \sum_{j=0}^{N+1} w_j^L (1 + x_j) v_N(x_j, t^m) v'_N(x_j, t^m) = \int_{-1}^1 w(x) (1 + x) v_N(x, t^m) v'_N(x, t^m) dx.$$

We integrate by parts the RHS of I, substitute $v_N(x, t^m) = (1 - x)p(x)$ from (3.35), and in view of (3.36) we obtain as before (compare (3.37)),

$$(6.17) \quad I \leq -2\eta_0 \|p\|_{(1-x)w(x)}^2.$$

The second expression, II, gives us

$$\begin{aligned} II &= \sum_{j=1}^N w_j a(x_j) (1 + x_j) [(1 - x_j) p'(x_j) - p(x_j)]^2 \leq \\ (6.18) \quad &\leq 2a_\infty \sum_{j=1}^N w_j (1 + x_j)^2 (p'(x_j))^2 + 2 \sum_{j=1}^N w_j a(x_j) (1 + x_j) p^2(x_j) \\ &= 2a_\infty II_1 + 2 \cdot II_2. \end{aligned}$$

The inverse inequality (2.6) with weight $\omega(x) = (1 - x)w(x)$ implies

$$II_1 = \|p'\|_{(1-x^2)(1-x)w(x)}^2 \leq \lambda_{N-1} \|p\|_{(1-x)w(x)}^2, \quad \lambda_{N-1} = \lambda_{N-1}(\alpha + 1, \beta)$$

and the expression II_2 does not exceed

$$II_2 \leq \max_{1 \leq j \leq N} [a(x_j) \frac{1 + x_j}{1 - x_j}] \cdot \sum_{j=0}^{N+1} w_j (1 - x_j) p^2(x_j) \leq 2 \cdot \max_{1 \leq j \leq N} \frac{a(x_j)}{1 - x_j} \cdot \|p\|_{(1-x)w(x)}^2.$$

Consequently we have

$$(6.19) \quad II \leq 2 \left(a_\infty \lambda_{N-1} + 2 \cdot \max_{1 \leq j \leq N} \frac{a(x_j)}{1 - x_j} \right) \|p\|_{(1-x)w(x)}^2.$$

Equipped with (6.17) and (6.18) we return to (6.16) to find

$$(6.20) \quad \|v_N(\cdot, t^{m+1})\|_\omega^2 \leq \|v_N(\cdot, t^m)\|_\omega^2 - 2\Delta t \left[2\eta_0 - \Delta t \left(a_\infty \lambda_{N-1} + 2 \max_{1 \leq j \leq N} \frac{a(x_j)}{1 - x_j} \right) \right] \|p\|_{(1-x)w(x)}^2,$$

and (6.18b) follows in view of the CFL condition (6.14b). \square

Notes.

1. The case $a(x_j) \equiv a = \text{Const.} > 0$, corresponds to the stability statement of Lemma 3.5. Similar stability statements with the appropriate weights which correspond to Lemmata

3.2, 3.3, and 3.4, namely, $\omega_j = \frac{1+x_j}{1-x_j} \frac{w_j^G}{a(x_j)}$, and $\omega_j = (1+x_j) \frac{w_j^G}{a(x_j)}$, hold. These statements cover the stability of the corresponding spectral and pseudospectral Jacobi approximations with variable coefficients.

2. We should highlight the fact the stability assertion stated in Theorem 6.2 depends solely on the uniform bound of $a(x_j)$ but otherwise is independent of the smoothness of $a(x)$.

3. The proof of Theorem 6.2 applies mutatis mutandis to the case of variable coefficients with $a = a(x, t)$. If $a(x_j, t)$ are C^1 -functions in the time variable then (6.20) is replaced by

$$\|v_N(\cdot, t^{m+1})\|_{\omega^{m+1}} \leq (1 + \text{Const.} \Delta t) \|v(\cdot, t^m)\|_{\omega^m}, \quad \omega_j^m = (1 + x_j) \frac{w_j^L}{a(x_j, t^m)}$$

and stability follows.

4. We conclude by noting that the CFL condition (6.14b) depends on the quantity $\max_{1 \leq j \leq N} \frac{a(x_j)}{1-x_j}$, rather than the minimal gridsize, $\frac{1}{\Delta x_{\min}}$, as in the constant coefficient case (compare (3.12)). This amplifies our introductory remarks in the beginning of this section, which claim that the $O(N^{-2})$ stability restriction is essentially due to the size of the Sturm-Liouville eigenvalues, $\lambda_{N-1} = O(N^2)$. Indeed, the other portion of the CFL condition requiring

$$\Delta t \cdot 2 \max_{1 \leq j \leq N} \frac{a(x_j)}{1-x_j} \leq \eta_0$$

guarantees the resolution of waves entering through the inflow boundary $x = 1$. In the constant coefficients case this resolution requires time-steps Δt of size $\frac{1}{\Delta x_{\min}}$. However, when the inflow boundary is almost characteristic, i.e., when $a(1) \sim 0$, then the CFL condition is essentially independent of Δx_{\min} , for (6.21) boils down to $\Delta t \cdot 2a'(1) \leq \eta_0$. In purely outflow cases such as (6.3), the time-step is independent of any resolution requirement at the boundaries and we are left with the CFL condition (6.5) stated in Theorem 6.1.

References

- [1] M. Abramowitz and I. A. Steger, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Government Printing Office, Washington, D. C., 1972.
- [2] C. Canuto, M. Y. Hussaini, A. Quarteroni, and T. Zang, *Spectral Methods in Fluid Dynamics*, Springer-Verlag, New York, 1988.
- [3] C. Canuto and A. Quarteroni, "Approximations results for orthogonal polynomials in Sobolev spaces," *Math. Comp.*, 38 (1982), pp. 67-86.
- [4] P. J. Davis and P. Rabinowitz, *Methods of Numerical Integration*, Academic Press, New York, 1975.
- [5] D. Gottlieb, "The stability of pseudospectral Chebyshev methods," *Math. Comp.*, 36 (1981), pp.107-118.
- [6] D. Gottlieb, L. Lustman, and E. Tadmor, "Stability analysis of spectral methods for hyperbolic initial-boundary value systems," *SINUM*, 24 (1987), pp. 241-256.
- [7] D. Gottlieb, L. Lustman, and E. Tadmor, "Convergence of spectral methods for hyperbolic initial-boundary value systems," *SINUM*, 24 (1987), pp. 532-537.
- [8] D. Gottlieb and S. Orszag, *Numerical Analysis of Spectral Methods: Theory and Applications*, SIAM, Philadelphia, PA, 1977.
- [9] B. Gustafsson, H. O. Kreiss, and A. Sundström, "Stability theory of difference approximations for mixed initial boundary value problems II," *Math. Comp.*, 26 (1972), pp. 649-688.
- [10] C. W. Shu, "Total-variation-diminishing time discretizations," *SISSC*, 6 (1988), pp. 1073-1084.
- [11] C. W. Shu and S. Osher, "Efficient implementation of essentially non-oscillatory shock-capturing schemes," *J. Comp. Phys.*, 77 (1988), pp. 439-471.
- [12] A. H. Stroud and D. Secrest, *Gaussian Quadrature Formulas*, Prentice-Hall, 1966.
- [13] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc., New York, 1967.

1. Report No. NASA CR-182053 ICASE Report No. 90-42		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle THE CFL CONDITION FOR SPECTRAL APPROXIMATIONS TO HYPERBOLIC INITIAL-BOUNDARY VALUE PROBLEMS				5. Report Date June 1990	
				6. Performing Organization Code	
7. Author(s) David Gottlieb Eitan Tadmor				8. Performing Organization Report No. 90-42	
				10. Work Unit No. 505-90-21-01	
9. Performing Organization Name and Address Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23665-5225				11. Contract or Grant No. NAS1-18605	
				13. Type of Report and Period Covered Contractor Report	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Langley Research Center Hampton, VA 23665-5225				14. Sponsoring Agency Code	
15. Supplementary Notes Langley Technical Monitor: Richard W. Barnwell To appear in Mathematics of Computation Cor = De Ma Final Report					
16. Abstract We study the stability of spectral approximations to scalar hyperbolic initial-boundary value problems with variable coefficients. Time is discretized by explicit multi-level or Runge-Kutta methods of order 3 (forward Euler time differencing is included), and we study spatial discretizations by spectral and pseudospectral approximations associated with the general family of Jacobi polynomials. We prove that these fully explicit spectral approximations are stable provided their time-step, Δt , is restricted by the CFL-like condition, $\Delta t < \text{Const. } N^{-2}$, where N equals the spatial number of degrees of freedom. We give two independent proofs of this result, depending on two different choices of appropriate L^2 -weighted norms. In both approaches, the proofs hinge on a certain inverse inequality interesting for its own sake. Our result confirms the commonly held belief that the above CFL stability restriction, which is extensively used in practical implementations, guarantees the stability (and hence the convergence) of fully-explicit/spectral approximations in the non-periodic case. <i>Keywords:</i> L-59 V/N-52					
17. Key Words (Suggested by Author(s)) hyperbolic equations, spectral approximations, stability condition. <i>G. J. G. 10</i>			18. Distribution Statement 64 - Numerical Analysis Unclassified - Unlimited		
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of pages 27	22. Price A03		